

let, also tibia 1, the tarsi, femur 1 save at basal half, knees, tibia 2 save widely at middle, and trochanters; hind knees black. Scaly, lower face somewhat rugulose. Propodeum with three sharp median and a lateral carina. Clothing sparse. Club 3-jointed, pedicel long, reddish, one half of club.

Deeral, jungle, July.

A NEW NOCTUID FROM OREGON

(*Lepidoptera, Noctuidæ, Acronyctinæ*)

BY HARRISON G. DYAR

Papaipema pertincta, new species.

Area beyond the curved outer line purplish, an ochreous mark before apex, the subterminal line showing as lightened denticulations, all as in *duovata* Bird, except that the color is smoother lustrous purplish; central area light ochreous near the margin, the area above median vein heavily dusted with dark brown; narrow basal area also clear, bounded by the double brown subbasal line; orbicular and claviform spots three, in an oblique line, pale yellow, not white, brown-ringed, the central one the smaller, the upper with an outwardly connected brown center; reniform dots four without, two within, and a central zigzag line, all pale yellow, distinctly larger than in *duovata* and not bicolored. Hind wing dark brownish gray, lighter in the cell, defining a large dark discal lunule. Expanse, 32 mm.

Type, male, No. 22833, U. S. Nat. Mus.; Forest Grove, Oregon, September 5, 1919, bred from *Lupinus polyphyllus* Lindl. (L. P. Rockwood).